

...and get Epidemiologic Reviews -
FREE

Mortality in a Long-term Open Cohort of Prostitute Women

John J. Potterat¹, Devon D. Brewer², Stephen Q. Muth¹, Richard B. Rothenberg³, Donald E. Woodhouse¹, John B. Muth¹, Heather K. Stites¹ and Stuart Brody⁴

+ Author Affiliations

Abstract

In this study, the authors estimated overall and cause-specific mortality among prostitute women. They recorded information on prostitute women identified by police and health department surveillance in Colorado Springs, Colorado, from 1967 to 1999. The authors assessed cause-specific mortality in this open cohort of 1,969 women using the Social Security Death Index and the National Death Index, augmented by individual investigations. They identified 117 definite or probable deaths and had sufficient information on 100 to calculate a crude mortality rate (CMR) of 391 per 100,000 (95% confidence interval (CI): 314, 471). In comparison with the general population, the standardized mortality ratio (SMR), adjusted for age and race, was 1.9 (95% CI: 1.5, 2.3). For the period of presumed active prostitution only, the CMR was 459 per 100,000 (95% CI: 246, 695) and the SMR was 5.9 (95% CI: 3.2, 9.0). Violence and drug use were the predominant causes of death, both during periods of prostitution and during the whole observation period. The CMR for death by homicide among active prostitutes was 229 per 100,000 (95% CI: 79, 378), and the SMR was 17.7 (95% CI: 6.2, 29.3). Deaths from acquired immunodeficiency syndrome occurred exclusively among prostitutes who admitted to injecting drug use or were inferred to have a history of it.

Key words [acquired immunodeficiency syndrome](#); [HIV](#); [homicide](#); [mortality](#); [overdose](#); [prostitution](#); [substance abuse](#), [intravenous](#); [violence](#)

Key words [Abbreviations: HIV, human immunodeficiency virus; NDI, National Death Index; SSDI, Social Security Death Index; SSN, Social Security number.](#)

Received for publication September 10, 2003; accepted for publication November 24, 2003.

Female prostitution is embedded in a context of felonious activity, illicit drugs, and violence (1-4) and is associated with premature mortality. Two cohort studies have directly assessed mortality in prostitutes during short observation periods (5, 6). No prior research has measured prostitute mortality during a long period or has verified it with vital statistics records. Indeed, Woolston's 1921 observation that "no satisfactory data showing the length of life of prostitutes in America are available, for the simple reason that women are not registered as such when they die" (7, p. 41) has remained accurate.

For three decades, continuous, community-wide surveillance of prostitutes in Colorado Springs, Colorado, has generated information on a large cohort of prostitute women (8, 9). These data and the availability of national mortality databases permitted assessment of cause-specific mortality among these women.

MATERIALS AND METHODS

Background and population characteristics

US Census figures (10) show that in 1970, Colorado Springs had a population of 235,972, of whom 94 percent were White (including an unknown proportion of Hispanics), 5 percent were African-American, and 1 percent were of other races. Thirty years later, the population of Colorado Springs had increased to 516,929 (11), of whom 81 percent were White (including 14 percent Hispanic), 7 percent were African-American, 4 percent were of Asian/Pacific Islander/American Indian race, and 9 percent were of other or mixed race (percentages do not sum to 100 because of rounding error).

Enhanced monitoring of prostitution in Colorado Springs began in the 1960s in response to a rapidly growing gonorrhea epidemic associated with prostitution (5, 8). In mid-1970, the health department instituted a mechanism for examining arrested prostitutes for sexually transmitted infections, termed the Health Hold Order (8). At about the same time, the police created a vice squad to monitor prostitution. Their activities included arrest and field surveillance (including the recording of personal identifiers) of persons engaged in prostitution. Colorado statutes urged coordination between public health officials and police officers in suppressing sexually transmitted infections and prostitution (5). This relationship was unilateral: Police provided health officers access to prostitution records but not vice versa. Between 1970 and 2000, health department and police prostitution records were periodically compared.

Health department records for prostitute women were also based on women identified independently in our sexually transmitted infection, human immunodeficiency virus (HIV), and drug treatment clinics (through self-report); in observations of prostitutes during public health outreach activities; and through contact tracing for sexually transmitted infections/HIV. In our clinics, we recorded standard information on each woman as part of routine clinical and epidemiologic evaluation. Starting in mid-1985, women visiting the sexually transmitted infection clinic or the HIV testing clinic who provided histories of prostitution were queried about injecting drug use and offered HIV testing; such information was unavailable for prostitutes ascertained solely by means of other sources.

The study cohort included women in Colorado Springs identified by police or health department surveillance as prostitutes, that is, as women who exchanged sex for money or drugs. These women were known to have engaged in prostitution between approximately 1967 and 1999. Most women ultimately appeared in multiple data sources during their prostitution careers (9). Both the prevalence and number of sexual partners of prostitute women in Colorado Springs appear to be representative of prostitutes in the United States (9, 12).

Mortality data sources

As a first step, we searched the Social Security Death Index (SSDI) in mid-1999 regarding people who had died between 1967 and 1999. The SSDI records information on decedents with Social Security numbers (SSNs) for whom survivor claims have been processed (13, 14). It lists the decedent's first name (at least first initial), surname, birth date, date of death, postal codes of last residence and last Social Security benefit, and SSN; it does not record information on sex, race, birth name (for women), death location, or cause of death. The staff of the sexually transmitted infection clinic requested the SSN as an optional item; 79 percent (981/1,245) of the women seen at the clinic provided it. Many (565/1,969) records of women in our cohort contained multiple names. Thus, we queried the SSDI systematically, beginning with SSN only and then using combinations of surname, first name/initial, and birth date to identify likely matches.

The National Death Index (NDI) registers deaths that have occurred in the United States since 1979. It is maintained by the Centers for Disease Control and Prevention. Source documentation consists of death certificate information provided by state and territorial registries. Minimum data required for searches

include first name and surname and either SSN or month/year of birth. Provision of middle initial, day of birth, father's surname, age at death, sex, race, marital status, state of residence, and/or state of birth improves search accuracy. NDI algorithms identify possible matches; the quality of matches depends on the completeness of submitted data.

Initial criteria for gauging the probability of a correct match are based on data concordance: 1) either first name or surname and SSN; 2) SSN and surname (for men) or SSN and surname or father's surname (for women); or 3) first name and surname and month and year of birth. The NDI program compensates for spelling subtleties by using phonetic codes. Possible matches include state of death and certificate number. The NDI search algorithm generates many potential matches, especially if the submitted information is incomplete, as well as a score (0–100). For all searches, we classified matches as *definite* (concordance of four of the following: SSN (within one digit), name, birth date, race, and sex); *probable* (among data items other than SSN, no more than two items slightly discordant (e.g., birth date one digit off or name slightly different)); and *possible* (manual review aiming for a preponderance of evidence using data from the NDI, the death certificate, and our records). We considered definite and probable deaths to be confirmed. Most NDI matches were eliminated using logic (e.g., males were automatically excluded) and a conservative NDI score cutoff point (less than 45 out of a possible 100 in the absence of an SSN).

Our list was searched at the NDI in late 1999 and compared with decedent records for 1979–1997. We sought death certificates from relevant registries in the United States for confirmed, probable, and possible matches emanating from the SSDI or NDI searches. Our NDI search was approved by the Colorado Health Sciences Center Institutional Review Board; in addition, death certificate requests were subject, in some states, to local institutional review. The Colorado Springs Police Department supplied a list of homicides that occurred in Colorado Springs between 1968 and 1999. We also requested information about death circumstances from other US jurisdictions for other murdered women in our cohort.

Statistical analyses

Mortality rates were based on the 1,633 women for whom year of first observation of local prostitution was available. Nearly all of the missing data on year of first local prostitution were attributable to women who did not have dated records in police files during the early part of the observation period, women identified only through visits to the HIV testing clinic or the drug treatment clinic (where dates of first local prostitution were not systematically collected), women who engaged in prostitution elsewhere, and women for whom the locale of prostitution could not be determined. Our calculations of person-years of observation took into account the fact that actual dates of first observation and death (when dates precise to the day and month were available) were uniformly distributed across a given year (resulting in 0.5 of a person-year, on average, for the first year a woman was observed or the year of her death) and that the final SSDI search occurred in mid-1999 (also resulting in 0.5 of a person-year for a woman censored by that endpoint).

Rates were computed for two sets of women: the cumulative cohort (which grew steadily from 1967 to 1999) and a subcohort classified as active prostitutes. The latter consisted of a shifting set of women. We created this "moving cohort" by including, for a given year, only women who were within 3 years of first being observed engaging in prostitution. Among prostitute women in our cohort known to reside in Colorado Springs for several years or more at some point prior to 1995 ($n = 449$), 50 percent had a time span between first and last observation of local prostitution (based on police surveillance, observation in the field or health department clinics, and/or reports from other active prostitute women) of 3 or more years. Fifteen percent

had a span of prostitution of 2 years, and 35 percent had a span of less than 2 years. Thus, we inferred that the majority of living women in the cohort continued to work as prostitutes while in the moving cohort.

We calculated the expected number of deaths (using the “prospective model” (15)) in the cumulative and active cohorts using sex-, age-, and race-specific mortality tables for all causes of death (16) and homicide (17-19). We computed standardized mortality ratios by dividing the observed number of deaths by the expected number of deaths. In addition, we estimated the average annual number of murdered prostitutes in the United States between 1981 and 1990 by multiplying the homicide rate we observed by the estimated number of prostitute women in the United States (23 per 100,000 population, derived from our local capture-recapture study (9)). Then we divided this product by the mean number of females, overall and between ages 15 and 44 years, murdered annually during this period (17-19) to estimate the percentage of female murder victims in the United States who were prostitutes. We estimated the typical prostitute’s risk of murder during her prostitution career by means of the formula $1 - (1 - \text{active prostitutes' crude homicide rate})^5$, because prostitutes in Colorado Springs worked for 5 years, on average (9).

RESULTS

Cohort characteristics

From 1967 through 1999, 1,969 women were identified as prostitutes in Colorado Springs. Nearly two thirds ($n = 1,245$; 63.2 percent) were initially identified in the sexually transmitted infection clinic, 134 (6.8 percent) were identified in the HIV testing center, 56 (2.8 percent) were identified in the substance abuse clinic, and 88 (4.5 percent) were identified during routine outreach. The remaining 446 women (22.7 percent) were initially identified by police; nearly three quarters (326/446) were observed by undercover officers, and the remainder were observed by uniformed officers patrolling solicitation areas. Four fifths (1,579/1,969) were known to have received care in one of our health department clinics. The overwhelming majority of the women worked as street prostitutes; only 126 worked in massage parlors, and most of these women also worked on the streets (9).

We classified the majority of prostitute women (1,795/1,969; 91 percent) as evanescent, short-term, or long-term residents. Evanescent prostitutes ($n = 1,012$; 56 percent) resided and solicited locally for a few days or weeks in any given year. Short-termers ($n = 315$; 18 percent) solicited for many weeks to months in any given year, and long-termers ($n = 468$; 26 percent) solicited for years even if they did not solicit continuously (9). Ninety-one percent were known to have engaged in prostitution locally, 4 percent reported histories of prostitution elsewhere only, and for 5 percent, locale was not recorded. The mean age at first observation of working locally was 24 years for those for whom we had data on date of first observation ($n = 1,633$; median, 23; standard deviation, 6.0; interquartile range, 20-27; range, 11-54).

Search results

The NDI search of 1,969 women used 2,883 names (mean = 1.47 per person; range, 1-9), yielding 9,926 potential matches, of which 1,223 met our initial matching criteria. Using the more rigorous matching criteria, we made 301 requests for death certificates for 165 women from 38 state registries. For the 33-year study interval, 117 women (6 percent) were classified as confirmed dead and 26 (1.3 percent) were classified as possibly dead (table 1). The 117 confirmed deaths were distributed in 26 states. Health department records and death certificates showed perfect concordance on SSN for 49 (86 percent) of the 57 confirmed-dead women who had an SSN in both sources. The NDI identified 100 confirmed deaths, with the SSDI corroborating only 57 of these; all deaths recorded by the SSDI appeared in the NDI. In reviewing the 43 deaths “missed” by the SSDI, at least 36 should have been listed, because a matching

name or SSN was available. Predictably (7), none of the death certificates recorded any history or evidence of prostitution.


View this table:
[In this window](#) [In a new window](#)

by race, 1967-1999

TABLE 1.
 Vital status of 1,969 women
 who worked as prostitutes in
 Colorado Springs, Colorado,


Comparative mortality

Observed dates of first prostitution were available for 1,633 women. Figures 1, 2, and 3 show for each year the number of women observed for the first time, the number in the cumulative cohort, and the number in the active cohort, respectively. Women who had dates of first prostitution were more likely to be African-American (33 percent) than women lacking such information (17 percent).




View larger version:
[In this page](#) [In a new window](#)
[Download as PowerPoint Slide](#)

FIGURE 1. Number of women
 observed for the first time
 working as prostitutes in
 Colorado Springs, Colorado,
 by year, 1967-1999.



View larger version:
[In this page](#) [In a new window](#)
[Download as PowerPoint Slide](#)

FIGURE 2. Number of
 prostitute women in the
 cumulative cohort, by year,
 Colorado Springs, Colorado,
 1967-1999.



View larger version:

FIGURE 3. Number of
 prostitute women in the
 active cohort, by year,
 Colorado Springs, Colorado,
 1967-1999.

[In this page](#) [In a new window](#)
[Download as PowerPoint Slide](#)

Of these 1,633 women, 100 died during the study period. The crude mortality rate was 391 per 100,000 person-years, and the standardized mortality ratio was 1.9 (table 2). The overall crude mortality rate and standardized mortality ratio for the presumed-active prostitutes—the women whose period of observation included only that time during which they were most likely to have been engaged in prostitution—were 459 per 100,000 person-years and 5.9, respectively.

View this table:

[In this window](#) [In a new window](#)

TABLE 2.
Overall mortality and
homicide rates among 1,969
prostitutes in Colorado

Springs, Colorado, 1967-1999

Causes of death

Few of the women died of natural causes, as would be expected for persons whose average age at death was 34 years. Rather, based on proportional mortality, the leading causes of death were homicide (19 percent), drug ingestion (18 percent), accidents (12 percent), and alcohol-related causes (9 percent) (table 3).

View this table:

[In this window](#) [In a new window](#)

TABLE 3.
Causes of death for
confirmed deaths among
1,969 prostitutes in Colorado

Springs, Colorado, 1967-1999

For nine women (8 percent), the underlying cause of death was HIV infection or acquired immunodeficiency syndrome; all had either admitted to a history of injecting drug use or were inferred (from having track marks, being named an injecting drug user by others, or associating only with other injecting drug users) to have a history of injecting drug use. Thus, deaths from acquired immunodeficiency syndrome occurred only among injecting drug users. Seven hundred seventy-four (98 percent) of the 786 women who had visited any of our health department clinics since mid-1985 had been tested for HIV, and 29 (3.7 percent) were HIV-positive. Twenty-seven of the 29 HIV-positive women admitted to injecting drug use. Among the 117 women who died, injecting drug use status was known for 59; 45 admitted to injecting drug use. Though incomplete, the data suggest that the nexus of injecting drug use, drug overdose, and acquisition of HIV was the most important nonviolent contributor to heightened mortality.

Violent death

Of 21 murders, nine occurred within 3 years of the first observed prostitution. All of these nine women were active prostitutes at the time of death, and eight were killed while soliciting. The crude mortality rate for homicide in the cumulative cohort was 82.0 per 100,000 person-years; for the cohort of active prostitutes, it was 229 per 100,000 person-years (table 2). On the basis of this latter estimate, these women faced, on average, a 1 percent (0.0114) chance of being murdered during their prostitution careers. By extrapolation, this rate implies that between 1981 and 1990, approximately 124 prostitute women were murdered annually in the United States, accounting for 2.5 percent of female murder victims (3.7 percent of female murder victims aged 15-44 years). The workplace homicide rate for prostitutes in the moving cohort (eight

murdered on the job) was 204 per 100,000 person-years. The standardized mortality ratio for homicide in the cumulative cohort was 7.9, and in the moving cohort it was 17.7. Thus, active prostitutes were almost 18 times more likely to be murdered than women of similar age and race during the study interval. The nine deaths occurring among active prostitutes that were not deemed to be homicides included five drug overdoses, two suicides, and two deaths due to unknown causes.

Although murder accounted for 19 percent of all confirmed deaths, it accounted for half of the 18 deaths in the active subcohort. Of the 12 women murdered more than 3 years after the first observation of prostitution, eight died while still actively working as prostitutes (six were slain while soliciting and another was killed by a boyfriend who was jealous of her prostitution activity). One victim was not actively engaged in prostitution; death circumstances suggested that two were killed by clients while soliciting; and no information on prostitution status was available for the other. Thus, the vast majority of murdered women in our sample were killed as a direct consequence of prostitution.

DISCUSSION

We collected data on 1,969 prostitute women in Colorado Springs. This large cohort, assembled over 30 years as part of routine sexually transmitted infection control efforts, afforded us the opportunity to generate approximate population-based mortality data on women who spent some time in "the life." Our analysis indicated that their high mortality was attributable primarily to violence and drug use. Notably, despite prostitutes' very high number of sexual partners (12), the only women in our cohort who died of acquired immunodeficiency syndrome had a history of injecting drug use. This latter result is consistent with the very strong association between injecting drug use and HIV infection among prostitute women in Israel, Vietnam, and elsewhere in the United States (20-22).

Our estimates of the all-cause crude mortality rate (459 per 100,000) and standardized mortality ratio (5.9) for presumed-active prostitutes are similar to earlier estimates from smaller groups of prostitute women followed in Colorado Springs in 1976-1977 (standardized mortality ratio = 3.3) (5), in Nairobi, Kenya, in 1998-2002 ($n = 466$; crude mortality rate = 310 per 100,000 (Stephen Moses, University of Manitoba, personal communication, 2003)), and in London, United Kingdom, in 1985-1994 ($n = 402$; crude mortality rate = 401 per 100,000; standardized mortality ratio = 12.2) (6). The London study was based on comparatively few (675) person-years, deaths were measured by incidental reports from other prostitutes, and the observation period for living women was the interval between the first and last clinic visits, truncating the observation period relative to women who died.

It is likely that we underestimated mortality in our current analysis. While the sensitivity of the SSDI relative to the NDI for the 1979-1997 period was only 57 percent, other research shows that the SSDI ascertained proportionally fewer deaths in the decades prior to 1979 than in the years after 1978 (23). Thus, the sensitivity of the SSDI for the 1967-1978 period in our study was probably lower than the sensitivity we estimated for the 1979-1997 period. Moreover, the NDI displayed only moderately high sensitivity (81-93 percent) for samples of known female decedents (24, 25). Had NDI data been available for the periods 1967-1978 and 1998-1999, for which we relied solely on the SSDI, we estimate that another 10 (9.7) additional confirmed deaths would have been identified. The extent of underestimation is probably even greater than that suggested by these factors, given that we classified deaths conservatively and did not include 26 possible deaths when computing mortality rates.

The relatively low sensitivity of the death indexes for our cohort might have been due to the tendency of women to have multiple surnames, to some

women's ineligibility for Social Security benefits (lack of sufficient legal employment), and to the dearth of information on fathers' surnames in our database. Other factors may have prevented ascertainment of death for some women. For example, prostitutes who died in a foreign country or were murdered but never identified as dead (because the body was never found or the remains were not identifiable) (26-28) would be unlikely to appear in death indexes. Some women also might have intentionally misreported identifying information to the health department or police.

To our knowledge, no population of women studied previously has had a crude mortality rate, standardized mortality ratio, or percentage of deaths due to murder even approximating those observed in our cohort. The workplace homicide rate for prostitutes (204 per 100,000) is many times higher than that for women and men in the standard occupations that had the highest workplace homicide rates in the United States during the 1980s (4 per 100,000 for female liquor store workers and 29 per 100,000 for male taxicab drivers) (29).

Our crude homicide mortality rate for presumed-active prostitutes (229 per 100,000) is also similar to the mortality rates extrapolated from passive and informal surveillance of prostitute women in Canada between 1992 and 1998 (30) (181 per 100,000 on the basis of our prostitute prevalence estimate (9) and national population figures (31)) and in the Canadian province of British Columbia between 1985 and 1990 (32) (112-225 per 100,000). Parallel calculations for the 34 known prostitute women who were murdered on the job in Canada between 1992 and 1995 (30) yield a workplace homicide rate of 127 per 100,000. In our study, murder accounted for 50 percent of the deaths among presumed-active prostitutes. Murder accounted for 29-100 percent of prostitute deaths observed in recent decades in Birmingham, United Kingdom (Hilary Kinnell, United Kingdom Network of Sex Work Projects, personal communication, 1999), Nairobi (Stephen Moses, University of Manitoba, personal communication, 2003), Vancouver (32), and London (6). However, prostitutes represented a greater share of all female murder victims in British Columbia between 1981 and 1990 (8 percent) (32) and in Canada overall between 1991 and 1995 (5 percent) (33) than in the United States between 1981 and 1990, by our estimate (3 percent).

The high homicide and overall mortality rates observed in our cohort probably reflect circumstances for nearly all prostitutes in the United States (where prostitution is illegal, except for a few rural Nevada counties where brothels are permitted (34)) and many other countries. Although these Colorado Springs prostitutes appeared to be representative of all US prostitutes in terms of prevalence and number of sexual partners (9, 12) and although they worked as prostitutes (and died) in many parts of the country, prostitutes elsewhere might have different mortality rates and profiles.

Clients perpetrate a large proportion of the lethal and nonlethal violence experienced by prostitutes (1, 32, 35-41). For instance, clients were suspected perpetrators in 55 (64 percent) of the 86 murders of known prostitute women that took place in Canada between 1992 and 1998 (30). Moreover, clients who are serial murderers may account for a disproportionate fraction of prostitute murders (26-28, 42-44). Surveys also indicate that prostitute women encounter more violence from clients when working on the streets than in off-street contexts (32, 45-47), and 84 percent of known prostitutes murdered in the United Kingdom in the 1990s worked on the streets (39).

Women engaged in prostitution face the most dangerous occupational environment in the United States. Research identifying individual and contextual factors that make prostitutes vulnerable to murder and drug overdose can inform the development of interventions for reducing harm (32, 37, 48, 49).

ACKNOWLEDGMENTS

This work was supported in part by grants R01 DA09928, R29 DA10640, U01 AI47473, and R01 MH58077 from the National Institutes of Health.

The authors thank Ken Basilio, formerly of the vital statistics office of the El Paso County Department of Health and Environment, for his exceptional tenacity in procuring death certificates.

Footnotes

 Correspondence to Dr. Devon D. Brewer, Interdisciplinary Scientific Research, P.O. Box 15110, Seattle, WA 98115 (website: <http://www.interscientific.net/contact.html>).

References

1. Farley M, Raval I, Kiramira M, et al. Prostitution in five countries: violence and post-traumatic stress disorder. *Fem Psychol* 1998;8:405-26.
[CrossRef](#)
2. Lowman J. Vancouver field study of prostitution research notes. (Working Papers on Pornography and Prostitution, report 8, vol 1). Ottawa, Ontario, Canada: Department of Justice Canada, 1984.
3. McKeganey N, Barnard M. Sex work on the streets: prostitutes and their clients. Buckingham, United Kingdom: Open University Press, 1996.
4. Stark CE. *Tricking and tripping: prostitution in the era of AIDS*. Putnam Valley, NY: Social Change Press, 2000.
5. Potterat JJ, Rothenberg PR, Bross DC. Gonorrhea in street prostitutes: epidemiologic and legal implications. *Sex Transm Dis* 1979;6:58-63.
[Medline](#) [Web of Science](#)
6. Ward H, Dav S, Weber J. Risky business: health and safety in the sex industry over a 9 year period. *Sex Transm Infect* 1999;75:340-3.
[Abstract](#)
7. Woolston HR. *Prostitution in the United States prior to the entrance of the United States into the World War*. Montclair, NJ: Patterson Smith, 1921.
8. Potterat JJ, Rothenberg PR, Muth IR, et al. Invoking, monitoring and relinquishing a public health police power: the Health Hold Order. *Sex Transm Dis* 1999;26:345-9.
[Medline](#) [Web of Science](#)
9. Potterat JJ, Woodhouse DE, Muth IR, et al. Estimating the prevalence and career longevity of prostitute women. *J Sex Res* 1990;27:233-43.
10. Bureau of the Census. *US Department of Commerce. 1970 census of the population*. Washington, DC: Bureau of the Census, 1971.
11. Bureau of the Census. *US Department of Commerce. Your gateway to Census 2000*. (Website). Washington, DC: Bureau of the Census, 2003. (World Wide Web URL: <http://www.census.gov/main/www/cen2000.html>).
12. Brewer DD, Potterat JJ, Carrott SC, et al. Prostitution and the sex discrepancy in reported number of sex partners. *Proc Natl Acad Sci U S A* 2000;97:12385-8.
[Abstract/FREE Full Text](#)
13. Secco HD, Paffenbarger PS, Lee L-M. Comparison of National Death Index and World Wide Web death searches. *Am J Epidemiol* 2000;152:107-11.
[Abstract/FREE Full Text](#)
14. Ancestry.com. Search the Social Security Death Index. (Website). Provo, UT: MyFamily.com, Inc. 2003. (World Wide Web URL: <http://www.ancestry.com/search/rectype/vital/ssdi/>).
15. Hartz AI, Giefer EE, Hoffmann RG. A comparison of two methods for calculating expected mortality. *Stat Med* 1983;2:381-6.
[Medline](#)
16. National Center for Health Statistics. *Centers for Disease Control and Prevention. Life tables*. (Website). Atlanta, GA: Centers for Disease Control and Prevention, 2003. (World Wide Web URL: <http://www.cdc.gov/nchs/products/pubs/pubd/lftbls/life/1966.htm>).
17. National Center for Health Statistics. *Centers for Disease Control and Prevention. Vital Statistics of the United States*. (Website). (Data for 1967-1978). Atlanta, GA:

- Centers for Disease Control and Prevention, 2003. (World Wide Web URL: <http://www.cdc.gov/nchs/products/pubs/pubd/vsus/vsus.htm>).
18. Centers for Disease Control and Prevention. WONDER [Wide-ranging Online Data for Epidemiologic Research]. (Website). (Data for 1979 and 1980). Atlanta, GA: Centers for Disease Control and Prevention, 2003. (World Wide Web URL: <http://wonder.cdc.gov/>).
19. National Center for Injury Prevention and Control. Centers for Disease Control and Prevention. WISQARS [Web-based Injury Statistics Query and Reporting System]. (Website). (Data for 1981-1999). Atlanta, GA: Centers for Disease Control and Prevention, 2003. (World Wide Web URL: <http://www.cdc.gov/ncipc/wisqars>).
20. Darrow W. Centers for Disease Control Collaborative Group for the Study of HIV-1 in Selected Women. Prostitution, intravenous drug use, and HIV-1 in the United States. In: Platt R, ed. AIDS, drugs, and prostitution. London, United Kingdom: Tavistock/Routledge, 1990:18-40.
21. Modan B, Goldschmidt R, Rubinstein E, et al. Prevalence of HIV antibodies in transsexual and female prostitutes. *Am J Public Health* 1992;82:590-2.
[Abstract/FREE Full Text](#)
22. Tuan NA, Hien NT, Chi PK, et al. Intravenous drug use among street-based sex workers: a high-risk behavior for HIV transmission. *Sex Transm Dis* 2004;31:15-19.
[Medline](#) [Web of Science](#)
23. Schnorr TM, Steanland K. Identifying deaths before 1979 using the Social Security Administration Death Master File. *Epidemiology* 1997;8:321-3.
[CrossRef](#) [Medline](#) [Web of Science](#)
24. Curb ID, Ford CE, Pressel S, et al. Ascertainment of vital status through the National Death Index and the Social Security Administration. *Am J Epidemiol* 1985;121:754-66.
[Abstract/FREE Full Text](#)
25. LaVoie TA, Dial C, Torres M, et al. Vital status in the National Panel Survey of Black Americans: a test of the National Death Index among African Americans. *J Natl Med Assoc* 1996;88:501-5.
[Medline](#)
26. Merlin R, White J. Bad trick: the hunt for Spokane's serial killer. Spokane, WA: New Media Ventures, 2001.
27. Smith C, Guillen T. The search for the Green River Killer. New York, NY: Penguin Putnam, 1991.
28. Greene T. Bad date: the lost girls of Vancouver's low track. Toronto, Ontario, Canada: ECW Press, 2001.
29. Castillo DM, Jenkins EJ. Industries and occupations at high risk for work-related homicide. *J Occup Med* 1994;36:125-32.
[CrossRef](#) [Medline](#) [Web of Science](#)
30. Lowman J. Violence and the outlaw status of (street) prostitution in Canada. *Violence Against Women* 2000;6:987-1011.
[Abstract/FREE Full Text](#)
31. Statistics Canada. Search Canadian statistics. (Website). Ottawa, Ontario, Canada: Statistics Canada, 2003. (World Wide Web URL: <http://www.statcan.ca/english/Pgdb/>).
32. Lowman J, Fraser J. Violence against persons who prostitute: the experience in British Columbia. (Technical report TR1996-14e). Ottawa, Ontario, Canada: Department of Justice Canada, 1996.
33. Duchesne D. Street prostitution in Canada. *Jurimet Service Bulletin* 17, no. 2). Ottawa, Ontario, Canada: Canadian Center of Justice Statistics, 1997.
34. Albert A. Brothel: the Mustang Ranch and its women. New York, NY: Random House, 2001.
35. Barry K. The prostitution of sexuality. New York, NY: New York University Press, 1995.
36. Day S, Ward H. Violence towards female prostitutes: violence in sex work extends to more than risks from clients. (Letter). *BMJ* 2001;323:230.
[CrossRef](#) [Medline](#)
37. El-Raceel N, Witte SS, Wada T, et al. Correlates of partner violence among female street-based sex workers: substance abuse, history of childhood abuse, and HIV risks. *AIDS Patient Care STDs* 2001;15:41-51.
[CrossRef](#) [Medline](#) [Web of Science](#)
38. Farley M, Barkan H. Prostitution, violence against women, and posttraumatic stress disorder. *Women Health* 1998;27:37-49.
[CrossRef](#) [Medline](#) [Web of Science](#)
39. Kinnell H. Violence against sex workers. (Online letter). *BMI* 2001;322. (World

Wide Web URL: <http://bmj.bmjournals.com/cgi/eletters/322/7285/524#13693>).

40. Silbert MH, Pines AM. Victimization of street prostitutes. *Victimology* 1982;7:122-33.
41. Vanwesenbeeck I, de Graaf R, van Zessen C, et al. Professional HIV risk taking, levels of victimization, and well-being in female prostitutes in the Netherlands. *Arch Sex Behav* 1995;24:503-15.
[CrossRef](#) [Medline](#) [Web of Science](#)
42. Dudek JA. When silenced voices speak: an exploratory study of prostitute homicide. (Doctoral dissertation). Philadelphia, PA: MCP Hahnemann University, 2001.
43. Newton M. *The encyclopedia of serial killers*. New York, NY: Checkmark Books, 2000.
44. Sudden P. *The complete history of Jack the Ripper*. New York, NY: Carroll and Graf Publishers, 1994.
45. Church S, Henderson M, Barnard M, et al. Violence by clients towards female prostitutes in different work settings: questionnaire survey. *BMJ* 2001;322:524-5.
[FREE Full Text](#)
46. Plumridge L, Abel G. A "segmented" sex industry in New Zealand: sexual and personal safety of female sex workers. *Aust N Z J Public Health* 2001;25:78-83.
[Medline](#) [Web of Science](#)
47. Dutt PM, Warr DJ. Vulnerability on the streets: female sex workers and HIV risk. *AIDS Care* 1997;9:539-47.
[Medline](#) [Web of Science](#)
48. Bellie DJ. Reduction of AIDS risk among 41 heroin addicted female street prostitutes: effects of free methadone maintenance. *J Addict Dis* 1993;12:7-23.
[CrossRef](#) [Medline](#)
49. Plummer J, Potterat JJ, Muth SQ, et al. Providing support and assistance for low income or homeless women. *JAMA* 1996;276:1874-5.
[Abstract/FREE Full Text](#)

Articles citing this article

Utility of the National Death Index in Ascertaining Mortality in Acquired Immunodeficiency Syndrome Surveillance

Am J Epidemiol (2011) 0(2011): kwr034v1-kwr034

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Prostitutes as Victims of Serial Homicide: Trends and Case Characteristics, 1970-2009

Homicide Studies (2011) 15(1): 74-100

[Abstract](#) [Full Text \(PDF\)](#)

Alcohol Use Among Female Sex Workers and Male Clients: An Integrative Review of Global Literature

Alcohol Alcohol (2010) 45(2): 188-199

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

A study of men who pay for sex, based on the Norwegian National Sex Surveys

Scand J Public Health (2010) 38(2): 135-140

[Abstract](#) [Full Text \(PDF\)](#)

Lack of association between heterosexual lifetime number of sex partners and prevalent HIV infection: a crucial implication

Int J STD AIDS (2010) 21(1): 74-75

[Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Increased Mortality in Bulimia Nervosa and Other Eating Disorders

Am. J. Psychiatry (2009) 166(12): 1342-1346

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Overestimation of heterosexually attributed AIDS deaths is associated with immature psychological defence mechanisms and clitoral masturbation during penile-vaginal intercourse

Int J STD AIDS (2009) 20(12): 869-875

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Police violence and sexual risk among female and transvestite sex workers in Serbia: qualitative study

BMJ (2008) 337(jul30_2): a811

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

The Missing Missing: Toward a Quantification of Serial Murder Victimization in the United States

Homicide Studies (2007) 11(4): 319-339

[Abstract](#) [Full Text \(PDF\)](#)

Migration, Sexual Exploitation, and Women's Health: A Case Report From a Community Health Center

Violence Against Women (2007) 13(5): 486-497

[Abstract](#) [Full Text \(PDF\)](#)

Extreme cause-specific mortality in a cohort of adult prisoners--1988 to 2002: a data-linkage study

Int J Epidemiol (2007) 36(2): 310-316

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Payment for living organ donation should be legalised.

BMJ (2006) 333(7571): 746-748

[Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Sex worker health: San Francisco style

Sex. Transm. Infect. (2006) 82(5): 418-422

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

What happens to women who sell sex? Report of a unique occupational cohort

Sex. Transm. Infect. (2006) 82(5): 413-417

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Is an internal comparison better than using national data when estimating mortality in longitudinal studies?

J. Epidemiol. Community Health (2006) 60(9): 819-821

[Abstract](#) [Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Prostitution in Vancouver: Violence and the Colonization of First Nations Women

Transcultural Psychiatry (2005) 42(2): 242-271

[Abstract](#) [Full Text \(PDF\)](#)

"Bad for the Body, Bad for the Heart": Prostitution Harms Women Even if Legalized or Decriminalized

Violence Against Women (2004) 10(10): 1087-1125

[Abstract](#) [Full Text \(PDF\)](#)

Prostitution laws: health risks and hypocrisy


CMAJ (2004) 171(2): 109

[Full Text \(HTML\)](#) [Full Text \(PDF\)](#)

Les lois contre la prostitution : risque pour la sante et hypocrisie

CMAJ (2004) 171(2): 111

[Full Text \(HTML\)](#) [Full Text \(PDF\)](#)



The Journal of Infectious Diseases
NEW to Oxford Journals



Other Oxford University Press sites:

